

Early Permian (Asselian) orthoconic cephalopods from the Taishaku Limestone, Akiyoshi Belt, Southwest Japan

Shuji Niko*, Masayuki Ehiro**, Osamu Nishikawa*** and Isao Nishikawa****

*Department of Environmental Studies, Faculty of Integrated Arts and Sciences, Hiroshima University, Higashihiroshima 739-8521, Japan, **The Tohoku University Museum, Sendai 980-8578, Japan, ***Mineral Industry Museum, Graduate School of Technology and Resource Science, Akita University, Akita 010-8502, Japan, ****Otsu 1797-1, Yuki, Jinsekikogen-cho, Jinseki-gun, Hiroshima 720-1812, Japan

Abstract: Orthoconic cephalopods including two species of the orthocerid nautiloids, *Michelinoceras hasei* sp. nov. and *Bogoslovskya miharanoroensis* Niko and Ozawa, 1997, and a bactritoid, *Aktastioceras nishikawai* Niko, Nishida and Hamada, 1993, are described from the Lower Permian (Asselian) limestone in the Miharanoro area of Hiroshima Prefecture, Southwest Japan. Stratigraphically, the limestone belongs to the Uyamano Formation of the Taishaku Limestone. *Michelinoceras hasei* differs from comparable taxa in the possession of the surface grooves.

Introduction

Following previous two papers (Niko et al., 1993; Ehiro et al., 2014) the present contribution deals subsequent results of detailed investigations of the cephalopod material that collected from the Taishaku Limestone over the ten years in 1950's to 1960's and preliminary examined by Hayasaka and Nishikawa (1962; faunal composition was introduced by Koizumi, 1967a, b, 1975, as the personal communications from the authors). Here, we additionally describe orthoconic cephalopods. They are *Michelinoceras hasei* sp. nov., *Bogoslovskya miharanoroensis* Niko and Ozawa, 1997, and *Aktastioceras nishikawai* Niko, Nishida and Hamada, 1993.

Systematic Paleontology

Class Cephalopoda Cuvier, 1797

Subclass Nautiloidea Agassiz, 1847

Order Orthocerida Kuhn, 1940

Superfamily Orthoceratoidea M'Coy, 1844

Family Orthoceratidae M'Coy, 1844

Subfamily Michelinoceratinae Flower, 1945

Genus *Michelinoceras* Foerste, 1932

Type species.—*Orthoceras michelini* Barrande, 1866.

Michelinoceras hasei sp. nov.

Figures 1.1–1.5, 1.14

Diagnosis.—Species of *Michelinoceras* with 4° in angle of conch expansion and transverse grooves as surface

ornamentation; camerae relatively short indicating 1.3–2.0 in form ratios (maximum width/length); siphuncle central and narrow.

Description.—An incomplete phragmocone is available for study, that indicates longiconic orthocone with gradual conch expansion having approximately 4° in angle; its approximate diameter and length are 9 mm near apical end and 51 mm, respectively; cross sections of conch are circular; shell surface ornamented by transverse fine grooves. Septa deeply concaved; sutures are not observable in the specimen, but apparent sinus and obliqueness of sutures are not recognized in longitudinal sections; cameral length relatively short for the genus; form ratios (maximum width/length) of camera are 1.3–2.0; siphuncle is central in position and consists of orthochoanitic septal necks and cylindrical connecting rings; length of septal necks is long, 0.9–1.2 mm; diameter of siphuncle is narrow, 0.9–1.0 mm; ratios of siphuncular diameter per shell diameter are approximately 0.1; endosiphuncular and cameral deposits are not developed.

Material examined.—holotype, IGPS coll. cat. no. 111503.

Occurrence and age.—*Michelinoceras hasei* sp. nov. was discovered in massive limestone at locality 1 in Ehiro et al., (2014, fig. 1) in the Miharanoro area of Tojo-cho, on the eastern end of Shobara City, Hiroshima Prefecture, Southwest Japan. Stratigraphically, the fossil-bearing limestone belongs to the Uyamano Formation of the central facies in the Taishaku Limestone (Hase et al., 1974). The age of the associated ammonoid fauna with *M. hasei* is highly likely to be Asselian (early Early Permian; Ehiro et al., 2014).

Etymology.—The specific name honors the late Dr. Akira Hase in recognition of his contributions for geology of the Taishaku Limestone.

Discussion.—*Michelinoceras hasei* sp. nov. somewhat resembles *M. magnum* Shimansky (1968, p. 53, 54, pl. 1, figs. 1a, b, v, 2) from the upper Lower Carboniferous of southern Urals in its gross conch shape, relatively short camera for the genus, and narrow siphuncle. However, the Russian species lacks surface ornamentation. *Michelinoceras?* sp. (Niko and Ozawa, 1997, p. 48, figs. 2.14–2.16) from the *Triticites contractus* Zone (late Gzhelian to early Asselian; late Late Carboniferous to early Early Permian) of the Taishaku Limestone also differs from the new species in having the smooth shell surface.

Genus ***Bogoslovskya*** Zhuravleva, 1978

Type species.—*Bogoslovskya perspicua* Zhuravleva, 1978.

Bogoslovskya miharanoroensis Niko and Ozawa, 1997
Figures 1.6, 1.7

Bogoslovskya miharanoroensis Niko and Ozawa, 1997, p. 48, 50, figs 2.1–2.9.

Description.—Longiconic orthocones with gradual conch expansion and circular (to slightly compressed?) cross sections; apical end of a relatively well-preserved specimen (IGPS coll. cat. no. 111499) has 11 mm in approximate diameter; shell surface ornamented by distant ridges forming very narrow ribs, whose direction is oblique and towards apex on siphuncular side. Sutures straight and transverse; siphuncle submarginal in position.

Material examined.—IGPS coll. cat. nos. 111498–111500. In addition, two poorly preserved specimens, IGPS coll. cat. nos. 111501 and 111502, are questionably assigned to this species.

Occurrence and age.—Same as *Michelinoceras hasei* sp. nov.

Discussion.—Morphologies of the present specimens are consistent with those of *Bogoslovskya miharanoroensis* described by Niko and Ozawa (1997) from the *Triticites contractus* Zone of the Taishaku Limestone. This species

resembles *B. akiyoshiensis* Niko, Nishida and Kyuma (1995, p. 193, 195, figs. 1.1–1.14) from the Moscovian (Upper Carboniferous) strata of the Akiyoshi Limestone, but in shell surface of the latter species is ornamented by fine transverse lirae. Another Middle Carboniferous species of *Bogoslovskya* from Japan, *B. omiensis* Niko (2001, p. 115–118, figs. 1.1–1.3, 1.5–1.9), clearly differs from *B. miharanoroensis* by its strongly eccentric siphuncular position.

Subclass Bactritoidea Shimansky, 1951

Order Bactritida Shimansky, 1951

Family Parabactritidae Shimansky, 1951

Genus ***Aktastioceras*** Shimansky, 1948

Type species.—*Aktastioceras kruglovi* Shimansky, 1948.

Aktastioceras nishikawai Niko, Nishida and Hamada,
1993
Figures 1.8–1.13, 1.15

Orthoceras cfr. *adrivnense*[sic]; Koizumi, 1967a, p. 135; 1967b, p. 208.

Pseudorthoceras[sic] cfr. *adrianense* (Gemmellaro, 1890); Koizumi, 1975, p. 22.


Aktastioceras nishikawai Niko, Nishida and Hamada, 1993, p. 314, 315, figs. 2.1–2.8.

Description.—Longiconic orthocones with approximately 9° in angle of conch expansion; cross sections of conchs are circular at apical shell and laterally compressed in adoral one; shell surface is marked by very fine lirae that run directly transverse to conch axis. Septa indicate deep concavities; except for dorsal sinus, sutures are straight and transverse; cameral length moderate for the genus; form ratios (maximum width/length) of camera are 1.8–2.3; siphuncle situates dorsal margin with orthochoanitic septal necks; length of septal necks is short, approximately 1.0 mm; diameter of septal foramen is approximately 0.8 mm; no connecting ring preserved;

Material examined.—IGPS coll. cat. nos. 111504 and 111505.

Occurrence and age.—Same as *Michelinoceras hasei* sp. nov.

Figure 1. Orthoconic cephalopods from the Taishaku Limestone in the Miharanoro area of Tojo-cho, on the eastern end of Shobara City, Hiroshima Prefecture. **1–5, 14.** *Michelinoceras hasei* sp. nov., holotype, IGPS coll. cat. no. 111503: 1, side view, $\times 1.5$; 2, cross sectional view of polished apical end, $\times 1.5$; 3, thin longitudinal section, $\times 3$; 4, 5, partial enlargements of Figure 1.3 to show details of siphuncular structures, $\times 10$; 14, surface ornamentation of transverse grooves, $\times 5$. **6, 7.** *Bogoslovskya miharanoroensis* Niko and Ozawa, 1997: 6, IGPS coll. cat. no. 111500, lateral view, siphuncular side on right, $\times 2$; 7, IGPS coll. cat. no. 111499, septal view, arrow indicates siphuncle, $\times 2$. **8–13, 15.** *Aktastioceras nishikawai* Niko, Nishida and Hamada, 1993: 8, 11, IGPS coll. cat. no. 111505; 8, side view, $\times 1.5$; 11, partial enlargements of Figure 1.8 to show surface ornamentation of very fine transverse lirae, $\times 3$; 9, 10, 12, 13, 15, IGPS coll. cat. no. 111504; 9, lateral view, venter on right, $\times 1.5$; 10, cross sectional view of adoral end, venter down, $\times 1.5$; 12, dorsal view, arrow indicates siphuncle, $\times 3$; 13, partial enlargement of Figure 1.15 to show details of siphuncular structure, $\times 8$; 15, dorsoventral polished section, venter on right, $\times 2$.



Discussion.—Records of bactritoid cephalopods from Japan are rare and restricted in allochthonous limestone bodies in the Akiyoshi Belt, including Akiyoshi (Niko et al., 1991), Omi (Niko, 2001), and Taishaku (Niko et al., 1993; Niko and Ozawa, 1997; this report).

Based on an associated label, it seems that a specimen (IGPS coll. cat. no. 111505) of *Aktastioceras nishikawai* was assigned tentatively to an orthocerid genus, *Pseudorthoceras* Girty, 1911, by Dr. Ichiro Hayasaka and Mr. Isao Nishikawa. This unpublished result was introduced by Koizumi (1967a, b, 1975). These erroneous assignments are corrected herein.

Acknowledgements

The specimens described here had been kept by Dr. Tamio Nishida after the examination of Hayasaka and Nishikawa (1962). He kindly permits the authors to study them. Dr. Yasunari Shigeta is thanked for loan of the types of *Aktastioceras nishikawai*. We also thank Dr. Masayuki Fujikawa for his helpful comments on the manuscript.

References

- Agassiz, L., 1847, *An Introduction to the Study of Natural History, in a Series of Lectures Delivered in the Hall of the College of Physicians and Surgeons*. New York, 58 p.
- Barrande, J., 1866, *Système Silurien du Centre de la Bohême. liere Partie: Recherches Paléontologiques. Vol. II. Céphalopdes*. Privately published, Prague and Paris, pls. 108–244.
- Cuvier, G., 1797, *Tableau Elementary de l'Histoire Naturelle des Animaux*. Baudouin, Paris, 710 p. (reissued by Bruxelles, Culture et Civilisation, Paris, 1969)
- Ehiro, M., Nishikawa, O., and Nishikawa, I., 2014, Early Permian (Asselian) ammonoids from the Taishaku Limestone, Akiyoshi Belt, Southwest Japan. *Paleontological Research*, vol. 18, p. 51–63.
- Flower, R. H., 1945, Classification of Devonian nautiloids. *American Midland Naturalist*, vol. 33, p. 675–724.
- Foerste, A. F., 1932, Black River and other cephalopods from Minnesota, Wisconsin, Michigan, and Ontario (Part I). *Denison University Bulletin, Journal of the Scientific Laboratories*, vol. 27, p. 47–136, pls. 7–37.
- Gemmellaro, G. G., 1890, La Fauna dei calcari con *Fusulina* della valle del Fiume Sosio (nella Provincia di Palermo). *Giornale di Scienze Naturali ed Economiche*, vol. 20, p. 37–138, pls. 11–19.
- Girty, G. H., 1911, On some new genera and species of Pennsylvanian fossils from the Wewoka Formation of Oklahoma. *Annals of the New York Academy of Sciences*, vol. 21, p. 119–156.
- Hase, A., Okimura, Y., and Yokoyama, T., 1974, The Upper Paleozoic formations in and around Taishaku-dai, Chugoku Massif, Southwest Japan; with special reference to the sedimentary facies of limestone. *Geological Report of the Hiroshima University*, no. 19, p. 1–39, pls. 1–8. (in Japanese with English abstract)
- Hayasaka, I., and Nishikawa, I., 1962, On some Permian macrofossils from Hiroshima Prefecture, Japan (A preliminary note). *Fossils (Palaeontological Society of Japan)*, no. 6, p. 27. (in Japanese)
- Koizumi, H., 1967a, On the Palaeozoic Permian cephalopods from Japan [I]. *Chigakukenkkyu*, vol. 18, p. 130–135. (in Japanese)
- Koizumi, H., 1967b, On the Palaeozoic Permian cephalopods from Japan [II]. *Chigakukenkkyu*, vol. 18, p. 207–210. (in Japanese)
- Koizumi, H., 1975, *Paleozoic Cephalopods of Japan*. Teiseki Bunko, Chiba, 149 p. (in Japanese)
- Kuhn, O., 1940, *Paläozoologie in Tabellen*. Fischer, Jena, 50 p.
- M'Coy, F., 1844, *A Synopsis of the Characters of the Carboniferous Limestone Fossils of Ireland*. Privately published, 274 p. (reissued by Williams and Norgate, London, 1862)
- Niko, S., 2001, Middle Carboniferous orthoconic cephalopods from the Omi Limestone Group, Central Japan. *Paleontological Research*, vol. 5, p. 115–120.
- Niko, S., Nishida, T., and Hamada, T., 1993, *Aktastioceras nishikawai* n. sp., a first Permian bactritoid cephalopod from Japan. *Journal of Paleontology*, vol. 67, p. 314–316.
- Niko, S., Nishida, T., and Kyuma, Y., 1991, Middle Carboniferous Bactritoidea (Mollusca: Cephalopoda) from the Akiyoshi Limestone Group, Yamaguchi Prefecture (Molluscan Paleontology of the Akiyoshi Limestone Group-X). *Transaction and Proceedings of the Palaeontological Society of Japan, New Series*, no. 161, p. 714–719.
- Niko, S., Nishida, T., and Kyuma, Y., 1995, A new Carboniferous cephalopod *Bogoslovskya akiyoshiensis* from Southwest Japan. *Transaction and Proceedings of the Palaeontological Society of Japan, New Series*, no. 179, p. 193–195.
- Niko, S., and Ozawa, T., 1997, Late Gzhelian (Carboniferous) to early Asselian (Permian) non-ammonoid cephalopods from the Taishaku Limestone Group, Southwest Japan. *Paleontological Research*, vol. 1, p. 47–54.
- Shimansky, V. N., 1948, Nekotorye novye orthotserakony iz artinskikh otlozhenii yuzhnogo Urala [Some new orthoceracones from Artinskian deposits of the southern Urals]. *Doklady Akademii Nauk SSSR*, vol. 60, p. 119–121, pl. 1. (in Russian)
- Shimansky, V. N., 1951, K voprosu ob evolyutsii verkhnepaleozoishikh pryamkh golovonogikh [On the evolution of the Upper Paleozoic straight nautiloids]. *Doklady Akademii Nauk SSSR*, vol. 79, p. 867–870. (in Russian)
- Shimansky, V. N., 1968, Kamennougolniye Orthoceratida, Oncoceratida, Actinoceratida i Bactritida [Carboniferous Orthoceratida, Oncoceratida, Actinoceratida and Bactritida]. *Akademiia Nauk SSSR, Trudy Paleontologicheskogo Instituta*, vol. 117, p. 1–151, pls. 1–20. (in Russian)
- Zhuravleva, F. A., 1978, Devonnskiye orthoserody, nadotryad Orthoceratoidea [Devonian orthocerids, superorder Orthoceratoidea]. *Akademiia Nauk SSSR, Trudy Paleontologicheskogo Instituta*, vol. 168, 223 p. (in Russian)